

Introduction - The Dead Sea Fault

Shmuel Marco

The Department of Geophysics and Planetary Sciences, Tel Aviv University.

E-mail: shmulkm@tau.ac.il

The Dead Sea Fault (DSF) accommodates sinistral motion between the Arabia plate and the Sinai subplate since the Middle Miocene, ~20 Ma. The interpretation of 107 km left-lateral slip along the DSF is based on observations from four independent sources: regional plate tectonics, local geology, seismology, and geodesy. The regional tectonics shows that the Red Sea is an incipient ocean, where the Arabian plate has been breaking away from Africa since Late Oligocene-Early Miocene. This motion is transferred to the collision with Eurasia via sinistral shear along the DSF (Courillot et al., 1987; Freund, 1965; Garfunkel, 1981; Joffe and Garfunkel, 1987; Quennell, 1956). Local geology shows systematic offset of numerous pre-Miocene geologic features by a total of ~107 km (Bartov et al., 1980; Freund, 1965; Quennell, 1956) and fault geometry that indicates left-lateral motion (Garfunkel, 1981). Paleoseismic and archaeoseismic studies show sub-recent activity as sinistral offsets of natural and of manmade structures (e.g., Amit et al., 2002; Ellenblum et al., 1998; Klinger et al., 2000; Meghraoui et al., 2003; Niemi et al., 2001). Focal mechanisms of moderate-to-large earthquakes show sinistral motion along the DSF and generally are in agreement with the location of the active faults, based on geological data (Baer et al., 1999; Hofstetter et al., 2007; Klinger et al., 1999; Salamon et al., 1996). And finally, geodetic measurements are consistent and confirm the left-lateral slip as well as the slip rate from other palaeoseismic evidence of 4 ± 1 mm/yr (Le Beon et al., 2006; Le Beon et al., 2008; McClusky et al., 2003; Reilinger et al., 2006; Wdowinski et al., 2004). This rate, as well as uniform Gutenberg-Richter frequency-magnitude relation, indicate stable tectonic regime in the last 60 ka (Begin et al., 2005; Hamiel et al., 2008).

The complex geometry of the fault is apparent in pull-apart grabens, which are associated with releasing bends, and pressure ridges that formed where restraining bends occur. Garfunkel (1981) maintains that the pull-apart basins are all shorter the total lateral offset because they began to

form at a later stage, after some motion had already accrued. This view is supported by seismic surveys that reveal earlier buried basins, which are no longer active (Frieslander, 2000).

The pull-apart basins have acted like sediment traps. Studies of the Miocene to Recent clastic and evaporitic sediments as well as some magmatic sequences that accumulated in the basins have yielded a wealth of information and insight on the history of sedimentological conditions and processes (e.g., Bookman et al., 2004; Frostick and Reid, 1989; Klinger et al., 2003; Sneh, 1981, 1982; Tsatskin and Nadel, 2003), climate (e.g., Bartov et al., 2003; Begin et al., 1974; Frumkin et al., 1991; Stein, 2001), geomagnetic secular variation (Marco et al., 1998), seismicity and deformation (e.g., Agnon et al., 2006; Bartov and Sagy, 2004; El-Isa and Mustafa, 1986; Heifetz et al., 2005; Ken-Tor et al., 2001; Marco et al., 1996; Migowski et al., 2004), fauna and flora (Kislev et al., 1992), humans, and environment (e.g., Braun et al., 1991; Goren-Inbar and Belitzky, 1989; Goren-Inbar et al., 2000; Ron and Levi, 2001).

Several authors noted that the detailed shape of the DSF had changed through time (Garfunkel, 1981; Heimann and Ron, 1987, 1993; Marco, 2007; Rotstein et al., 1992; Shamir et al., 2005; ten Brink et al., 1999; ten-Brink and Ben-Avraham, 1989). The widest zone of about 50 km of distributed faulting is found in the Galilee, where the early-stage (Miocene) faults were associated with formation of basins (Freund et al., 1970; Shaliv, 1991) and with rotation of rigid blocks about sub-vertical axes (Ron et al., 1984). Subsequent post-Miocene deformation took place mostly in the form of normal faulting on E-W trending faults and the transform movement is currently localized in a very narrow zone. The deformation in the south was characterized initially by a 20-30-km-wide zone with primarily strike-slip and some normal slip on faults trending sub-parallel to the main transform fault. It later became localized in the Arava, where a single narrow fault zone offsets the youngest alluvium. In the earliest phase, young faults became active in the Negev, some 20 km west of the Arava (Avni et al., 2000), perhaps indicating another widening phase of the DSF zone (Marco, 2007).

Table 1. Various estimates of the Dead Sea Fault slip rate

Period	Rate mm/y	Data	Reference
Late Pleistocene-Recent	10	Geological	(Freund et al., 1968)
Last 1000 yr	0.8-1.7	Historical	(Garfunkel et al., 1981)

Plio-Pleistocene	7-10	Geological	(Garfunkel et al., 1981)
Last 4500 yr	2.2	Seismicity	(Ben-Menahem, 1981)
Late Pleistocene	6.4±0.4	Seismicity	(El-Isa and Mustafa, 1986)
Plio-Pleistocene	6 (0.283°/ma)	Plate kinematics	(Joffe and Garfunkel, 1987)
Holocene	9	Geological	(Reches and Hoexter, 1981)
Plio-Pleistocene	20	Geological	(Steinitz and Bartov, 1986)
Holocene	>0.7	Geological	(Gardosh et al., 1990)
Plio-Pleistocene	5.4-6.1	Geological	(Heimann, 1990)
Plio-Pleistocene	3-7	Drainage systems, Arava Fault	(Ginat et al., 1998)
Pleistocene	2-6, prefer 4	Alluvial fans, N. Arava	(Klinger et al., 2000)
Pleistocene	4.7±1.3	Alluvial fans, Arava	(Niemi et al., 2001)
Last 2000 yrs	6.9±0.1	Paleo and Archaeoseismology, Missyaf (DSF in Syria)	(Meghraoui et al., 2003)
1996-1999	2.6±1	Geodesy, GPS	(Pe'eri et al., 2002)
1996-2003	3.3±0.4	Geodesy, GPS	(Wdowinski et al., 2004)
25 ka	3.8-6.4	Geological, Lebanon	(Daëron et al., 2004)
Last 5000 yrs	≥3	Stream channel, Jordan Gorge	(Marco et al., 2005)
Survey-Mode GPS	5.6 to 7.5	(from south to north)	(McClusky et al., 2003)
1999-2005	4.9±1.4	GPS	(Le Beon et al., 2008) and thesis
Last 47.5 kyrs	4.7 to 5.1 mm/yr	Offset channels, Jordan	(Ferry et al., 2007) Comment/Reply: (Ferry and Meghraoui, 2008; Klein, 2008)

References

- Agnon, A., Migowski, C., and Marco, S., 2006, Intraclast breccia layers in laminated sequences: recorders of paleo-earthquakes, *in* Enzel, Y., Agnon, A., and Stein, M., eds., *New Frontiers in Dead Sea Paleoenvironmental Research*, Geological Society of America Special Publication, p. 195-214.
- Amit, R., Zilberman, E., Enzel, Y., and Porat, N., 2002, Paleoseismic evidence for time dependency of seismic response on a fault system in the southern Arava valley, Dead Sea rift, Israel: *Geol. Soc. Am. Bull.*, v. 114, p. 192-206.

- Avni, Y., Bartov, Y., Garfunkel, Z., and Ginat, H., 2000, Evolution of the Paran drainage basin and its relation to the Plio-Pleistocene history of the Arava Rift western margin, Israel: *Isr. J. Earth Sci.*, v. 49, p. 215-238.
- Baer, G., Sandwell, D., Williams, S., Bock, Y., and Shamir, G., 1999, Coseismic deformation associated with the November 1995, Mw = 7.1 Nuweiba earthquake, Gulf of Elat (Aqaba), detected by synthetic aperture radar interferometry: *J. Geophys. Res.*, v. 104, p. 25,221-25,232.
- Bartov, Y., Goldstein, S.L., Stein, M., and Enzel, Y., 2003, Catastrophic arid episodes in the Eastern Mediterranean linked with the North Atlantic Heinrich events: *Geology*, v. 31, p. 439-442.
- Bartov, Y., and Sagy, a., 2004, Late Pleistocene extension and strike-slip in the Dead Sea Basin: *Geological Magazine*, v. 141, p. 565-572.
- Bartov, Y., Steinitz, G., Eyal, M., and Eyal, Y., 1980, Sinistral movement along the Gulf of Aqaba - its age and relation to the opening of the Red Sea: *Nature*, v. 285, p. 220-221.
- Begin, B.Z., Steinberg, D.M., Ichinose, G.A., and Marco, S., 2005, A 40,000 years unchanging of the seismic regime in the Dead Sea rift: *Geology*, v. 33, p. 257-260.
- Begin, Z.B., Ehrlich, A., and Nathan, Y., 1974, Lake Lisan, the Pleistocene precursor of the Dead Sea: *Geol. Surv. Isr. Bull.*, v. 63, p. 30.
- Ben-Menahem, A., 1981, Variation of slip and creep along the Levant Rift over the past 4500 years: *Tectonophysics*, v. 80, p. 183-197.
- Bookman, R., Enzel, Y., Agnon, A., and Stein, M., 2004, Late Holocene lake levels of the Dead Sea: *Bull. Geol. Soc. Am.*, v. 116, p. 555-571.
- Braun, D., Ron, H., and Marco, S., 1991, Magnetostratigraphy of the hominid tool bearing Erkel Ahmar formation in the northern Dead Sea Rift: *Isr. J. Earth Sci.*, v. 40, p. 191-197.
- Courtillot, V., Armijo, R., and Tapponnier, P., 1987, The Sinai Triple Junction Revisited: *Tectonophysics*, v. 141, p. 181-190.

- Daëron, M., Benedetti, L., Tapponnier, P., Sursock, A., and Finkel, R.C., 2004, Constraints on the post 25-ka slip rate of the Yammouneh fault (Lebanon) using in situ cosmogenic ³⁶Cl dating of offset limestone-clast fans: *Earth Planet. Sci. Lett.*, v. 227, p. 105-119.
- El-Isa, Z.H., and Mustafa, H., 1986, Earthquake deformations in the Lisan deposits and seismotectonic implications: *Geophys. J. R. Astron. Soc.*, v. 86, p. 413-424.
- Ellenblum, R., Marco, S., Agnon, A., Rockwell, T., and Boas, A., 1998, Crusader castle torn apart by earthquake at dawn, 20 May 1202: *Geology*, v. 26, p. 303-306.
- Ferry, M., and Meghraoui, M., 2008, Reply to the comment of Dr M. Klein on: "A 48-kyr-long slip rate history for the Jordan Valley segment of the Dead Sea Fault": *Earth and Planetary Science Letters*, v. 268, p. 241-242.
- Ferry, M., Meghraoui, M., Abou Karaki, N., Al-Taj, M., Amoush, H., Al-Dhaisat, S., and Barjous, M., 2007, A 48-kyr-long slip rate history for the Jordan Valley segment of the Dead Sea Fault: *Earth and Planetary Science Letters*, v. 260, p. 394-406.
- Freund, R., 1965, A model of the structural development of Israel and adjacent areas since Upper Cretaceous times: *Geol. Mag.*, v. 102, p. 189-205.
- Freund, R., Garfunkel, Z., Zak, I., Goldberg, M., Weissbrod, T., and Derin, B., 1970, The shear along the Dead Sea rift: *Phil. Trans. Roy. Soc. Lond.*, v. A. 267, p. 107-130.
- Freund, R., Zak, I., and Garfunkel, Z., 1968, Age and rate of the sinistral movement along the Dead Sea Rift: *Nature*, v. 220, p. 253-255.
- Frieslander, U., 2000, The structure of the Dead Sea Transform emphasizing the Arava using new geophysical data [Ph.D. (in Hebrew, English abstract) thesis], The Hebrew University of Jerusalem.
- Frostick, L.E., and Reid, I., 1989, Climatic versus tectonic controls of fan sequences: lessons from the Dead Sea, Israel: *Jour. Geol. Soc. London*, v. 146, p. 527-538.
- Frumkin, A., Magaritz, M., Carmi, I., and Zak, I., 1991, The Holocene climatic record of the salt caves of Mount Sedom, Israel: *The Holocene*, v. 3, p. 191-200.

- Gardosh, M., Reches, Z., and Garfunkel, Z., 1990, Holocene tectonic deformation along the western margins of the Dead Sea: *Tectonophysics*, v. 180, p. 123-137.
- Garfunkel, Z., 1981, Internal structure of the Dead Sea leaky transform (rift) in relation to plate kinematics: *Tectonophysics*, v. 80, p. 81-108.
- Garfunkel, Z., Zak, I., and Freund, R., 1981, Active faulting in the Dead Sea rift: *Tectonophysics*, v. 80, p. 1-26.
- Ginat, H., Enzel, Y., and Avni, Y., 1998, Translocation of Plio-Pleistocene drainage system along the Dead Sea Transform, south Israel: *Tectonophysics*, v. 284, p. 151-160.
- Goren-Inbar, N., and Belitzky, S., 1989, Structural position of the Pleistocene Gesher Benot Ya'akov site in the Dead Sea Rift zone: *Quaternary Research*, v. 31, p. 371-376.
- Goren-Inbar, N., Craig, F.S., Verosub, K.L., Melamed, Y., Kislev, M., Tchernov, E., and Saragusti, I., 2000, Pleistocene milestones on the Out-of-Africa Corridor at Gesher Benot Ya'akov, Israel: *Science*, v. 289, p. 944-947.
- Hamiel, Y., Amit, R., Begin, Z.B., Marco, S., Katz, O., Salamon, A., Zilberman, E., and Porat, N., 2008, The seismicity along the Dead Sea fault during the last 60,000 years: *Seismological Society of America Bulletin*, v. in press.
- Heifetz, E., Agnon, A., and Marco, S., 2005, Soft sediment deformation by Kelvin Helmholtz Instability: A case from Dead Sea earthquakes: *Earth Planet. Sci. Lett.*, v. 236, p. 497-504.
- Heimann, A., 1990, The development of the Dead Sea rift and its margins in the northern Israel during the Pliocene and the Pleistocene, *Golan Res. Inst. and Geol. Surv. Isr.*
- Heimann, A., and Ron, H., 1987, Young faults in the Hula pull-apart basin, central Dead Sea transform: *Tectonophysics*, v. 141, p. 117-124.
- , 1993, Geometric changes of plate boundaries along part of the northern Dead Sea Transform: Geochronologic and paleomagnetic evidence: *Tectonics*, v. 12, p. 477-491.

- Hofstetter, R., Klinger, Y., Amrat, A.Q., Rivera, L., and Dorbath, L., 2007, Stress tensor and focal mechanisms along the Dead Sea fault and related structural elements based on seismological data: *Tectonophysics*, v. 429, p. 165-181.
- Joffe, S., and Garfunkel, Z., 1987, Plate kinematics of the circum Red Sea- a re-evaluation: *Tectonophysics*, v. 141, p. 5-22.
- Ken-Tor, R., Agnon, A., Enzel, Y., Marco, S., Negendank, J.F.W., and Stein, M., 2001, High-resolution geological record of historic earthquakes in the Dead Sea basin: *J. Geophys. Res.*, v. 106, p. 2221-2234.
- Kislev, M.E., Nadel, D., and Carmi, I., 1992, Epipalaeolithic (19,000 Bp) Cereal and Fruit Diet at Ohalo-Ii, Sea of Galilee, Israel: *Review of Palaeobotany and Palynology*, v. 73, p. 161-166.
- Klein, M., 2008, A comment on: "A 48-kyr-long slip rate history for the Jordan Valley segment of the Dead Sea Fault" *EPSL* 260 (2007) 394-406: *Earth and Planetary Science Letters*, v. 268, p. 239-240.
- Klinger, Y., Avouac, J.P., Abou-Karaki, N., Dorbath, L., Bourles, D., and Reyss, J.L., 2000, Slip rate on the Dead Sea transform fault in northern Araba Valley (Jordan): *Geophys. J. Int.*, v. 142, p. 755-768.
- Klinger, Y., Avouac, J.P., Bourles, D., and Tisnerat, N., 2003, Alluvial deposition and lake-level fluctuations forced by Late Quaternary climate change: the Dead Sea case example: *Sedimentary Geology*, v. 162, p. 119-139.
- Klinger, Y., Rivera, L., Haessler, H., and Maurin, J.C., 1999, Active faulting in the Gulf of Aqaba: New knowledge from the Mw7.3 earthquake of 22 November 1995: *Bull. Seism. Soc. Am.*, v. 89, p. 1025-1036.
- Le Beon, M., Klinger, Y., Agnon, A., Dorbath, L., Baer, G., Meriaux, A.-S., Ruegg, J.-C., Charade, O., Finkel, R., and Ryerson, F., 2006, Geodetic versus geologic slip rate along the Dead Sea, *Seismological Society of America Annual Meeting*: San Francisco.

- Le Beon, M., Klinger, Y., Amrat, A.Q., Agnon, A., Dorbath, L., Baer, G., Ruegg, J.C., Charade, O., and Mayyas, O., 2008, Slip rate and locking depth from GPS profiles across the southern Dead Sea Transform: *Journal of Geophysical Research-Solid Earth*, v. 113, p. -.
- Marco, S., 2007, Temporal variation in the geometry of a strike-slip fault zone: Examples from the Dead Sea Transform: *Tectonophysics*, v. 445, p. 186 – 199.
- Marco, S., Rockwell, T.K., Heimann, A., Frieslander, U., and Agnon, A., 2005, Late Holocene slip of the Dead Sea Transform revealed in 3D palaeoseismic trenches on the Jordan Gorge segment: *Earth Planet. Sci. Lett.*, v. 234, p. 189-205.
- Marco, S., Ron, H., McWilliams, M.O., and Stein, M., 1998, High-Resolution Record of Geomagnetic Secular Variation from Late Pleistocene Lake Lisan Sediments (Paleo Dead Sea): *Earth Planet. Sci. Lett.*, v. 161, p. 145-160.
- Marco, S., Stein, M., Agnon, A., and Ron, H., 1996, Long term earthquake clustering: a 50,000 year paleoseismic record in the Dead Sea Graben: *J. Geophys. Res.*, v. 101, p. 6179-6192.
- McClusky, S., Reilinger, R., Mahmoud, S., Sari, D.B., and Tealeb, A., 2003, GPS constraints on Africa (Nubia) and Arabia plate motions: *Geophys. J. Int.*, v. 155, p. 126.
- Meghraoui, M., Gomez, F., Sbeinati, R., derWoerd, J.V., Mouty, M., Darkal, A.N., Radwan, Y., Layyous, I., Najjar, H.A., Darawcheh, R., Hijazi, F., Al-Ghazzi, R., and Barazangi, M., 2003, Evidence for 830 years of seismic quiescence from palaeoseismology, archaeoseismology and historical seismicity along the Dead Sea fault in Syria: *Earth Planet. Sci. Lett.*, v. 210, p. 35-52.
- Migowski, C., Agnon, A., Bookman, R., Negendank, J.F.W., and Stein, M., 2004, Recurrence pattern of Holocene earthquakes along the Dead Sea transform revealed by varve-counting and radiocarbon dating of lacustrine sediments: *Earth and Planetary Science Letters*, v. 222, p. 301-314.
- Niemi, T.M., Zhang, H., Atallah, M., and Harrison, B.J., 2001, Late Pleistocene and Holocene slip rate of the Northern Wadi Araba fault, Dead Sea Transform, Jordan: *Journal of Seismology*, v. 5, p. 449-474.

- Pe'eri, S., Wdowinski, S., Shtibelman, A., and Bechor, N., 2002, Current plate motion across the Dead Sea Fault from three years of continuous GPS monitoring: *Geophys. Res. Lett.*, v. 29, p. doi: 10.1029/2001GL013879.
- Quennell, A.M., 1956, Tectonics of the Dead Sea rift, Congreso Geologico Internacional, 20th session, Asociacion de Servicios Geologicos Africanos: Mexico City, p. 385-405.
- Reches, Z., and Hoexter, D.F., 1981, Holocene seismic and tectonic activity in the Dead Sea area: *Tectonophysics*, v. 80, p. 235-254.
- Reilinger, R., McClusky, S., Vernant, P., Lawrence, S., Ergintav, S., Cakmak, R., Ozener, H., Kadirov, F., Guliev, I., Stepanyan, R., Nadariya, M., Hahubia, G., Mahmoud, S., Sakr, K., ArRajehi, A., Paradissis, D., Al-Aydrus, A., Prilepin, M., Guseva, T., Evren, E., Dmitrotsa, A., Filikov, S.V., Gomez, F., Al-Ghazzi, R., and Karam, G., 2006, GPS constraints on continental deformation in the Africa-Arabia-Eurasia continental collision zone and implications for the dynamics of plate interactions: *Journal of Geophysical Research-Solid Earth*, v. 111, p. -.
- Ron, H., Freund, R., Garfunkel, Z., and Nur, A., 1984, Block-rotation by strike-slip faulting: structural and paleomagnetic evidence: *J. Geophys. Res.*, v. 89, p. 6256-6270.
- Ron, H., and Levi, S., 2001, When did hominids first leave Africa?: New high-resolution magnetostratigraphy from the Erk-el-Ahmar Formation, Israel: *Geology*, v. 29, p. 887-890.
- Rotstein, Y., Bartov, Y., and Frieslander, U., 1992, Evidence for local shifting of the main fault and changes in the structural setting, Kinarot basin, Dead Sea transform: *Geology*, v. 20, p. 251-254.
- Salamon, A., Hofstetter, A., Garfunkel, Z., and Ron, H., 1996, Seismicity of the eastern Mediterranean region: Perspective from the Sinai subplate: *Tectonophysics*, v. 263, p. 293-305.
- Shaliv, G., 1991, Stages in the tectonic and volcanic history of the Neogene basin in the Lower Galilee and the valleys: Geological Survey of Israel - Report (in Hebrew, English abstract), v. 11/91, p. 1-94.

- Shamir, G., Eyal, Y., and Bruner, I., 2005, Localized versus distributed shear in transform plate boundary zones: The case of the Dead Sea Transform in the Jericho Valley: *Geochemistry Geophysics Geosystems*, v. 6, p. -.
- Sneh, A., 1981, The Hazeva Formation in the northern Arava: *Isr. Jour. Earth Sci.*, v. 30, p. 81-92.
- , 1982, Quaternary of the northwestern Arava, Israel: *Isr. Jour. Earth Sci.*, v. 31, p. 9-16.
- Stein, M., 2001, The sedimentary and geochemical record of Neogene - Quaternary water bodies in the Dead Sea Basin - inferences for the regional paleoclimatic history: *Journal of Paleolimnology*, v. 26, p. 271-282.
- Steinitz, G., and Bartov, Y., 1986, The 1985 time table for the tectonic events along the Dead Sea transform: *Terra Cognita*, v. 6, p. 160.
- ten Brink, U.S., Rybakov, M., Al-Zoubi, A.S., Hassouneh, M., Frieslander, U., Batayneh, A.T., Goldschmidt, V., Daoud, M.N., Rotstein, Y., and Hall, J.K., 1999, Anatomy of the Dead Sea transform: Does it reflect continuous changes in plate motion?: *Geology*, v. 27, p. 887-890.
- ten-Brink, U.S., and Ben-Avraham, Z., 1989, The anatomy of a pull-apart basin: Reflection observations of the Dead Sea basin: *Tectonics*, v. 8, p. 333-350.
- Tsatskin, A., and Nadel, D., 2003, Formation processes at the Ohalo II submerged prehistoric campsite, Israel, inferred from soil micromorphology and magnetic susceptibility studies: *Geoarchaeology-an International Journal*, v. 18, p. 409-432.
- Wdowinski, S., Bock, Y., Baer, G., Prawirodirdjo, L., Bechor, N., Naaman, S., Knafo, R., Forrai, Y., and Melzer, Y., 2004, GPS Measurements of current crustal movements along the Dead Sea Fault: *J. Geophys. Res.*, v. 109, p. 1-16.